

**CAUTION**  
**KEEP OUT OF REACH OF CHILDREN**  
**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

# **Terrazole<sup>®</sup> 350WP**

**SOIL FUNGICIDE**

**ACTIVE CONSTITUENT: 350 g/kg ETRIDIAZOLE**

**GROUP 14 FUNGICIDE**

**Controls damping off, root & stem rot of ornamentals & turf caused by *Phytophthora* & *Pythium* fungi as specified in the DIRECTIONS FOR USE table**

**Net Contents: 10 kg**

Arysta LifeScience Australia Pty Ltd  
Level 7, 435 King William Street Adelaide SA 5000  
Tel: (08) 8112 0900 Fax: (08) 8112 0999

09/15/F



## DIRECTIONS FOR USE

Crop	Disease	States	Rate	Critical Comments
<b>BEDDING PLANTS (ornamental seedlings)</b>	Root Rot, Stem Rot ( <i>Phytophthora</i> spp) Damping Off ( <i>Pythium</i> spp)	ALL STATES	60-100 g in 250 L water to 40 m <sup>2</sup>	Apply as a drench. Repeat applications may be made at 4 - 6 week intervals if required. Use higher rate on soils high in organic matter.
<b>HERBACEOUS AND WOODY ORNAMENTAL PLANTS</b>			60-100 g in 100 L water to 10 m <sup>2</sup>	Repeat applications at 2-3 month intervals to maintain disease control.
<b>BEDS POTS</b>			25-75 g in 100 L water	Apply 300 mL of solution per 15 cm pot or sufficient solution to saturate the soil in the pot. Use higher rate for soils with high organic matter or peat.
<b>TURF ESTABLISHED</b>	Damping Off Grease Spot ( <i>Pythium</i> spp)	NSW, ACT, Vic, SA and WA	100-250 g in 25 L water to 100 m <sup>2</sup>	Apply uniformly as a spray. Repeat treatment at 5-10 days to maintain disease control.
<b>TURF NEWLY SEEDED</b>			150-200 g in 25-50 L water to 100 m <sup>2</sup>	Apply as a spray immediately after seeding. Repeat application at 5-7 day intervals to maintain disease control.
<b>DRY SOIL MIX (seed box and potting mixes)</b>	Damping Off ( <i>Pythium</i> spp)	ALL STATES	45-90 g per m <sup>3</sup> of soil	Apply as dust and mix into one cubic metre of soil during soil fertilizer blending operation.
<b>ROOT DIP</b>	Root infections ( <i>Phytophthora</i> spp)	Qld, NSW, ACT, WA, Tas and SA	60-70 g in 50 L water	Mix well and stir frequently to maintain adequate suspension during dipping operation.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**WITHHOLDING PERIOD:  
NOT REQUIRED WHEN USED AS DIRECTED.**

## GENERAL INSTRUCTIONS

### Mixing

Add the required quantity to the water in the spray tank with agitation. Spray solutions need to be continuously agitated after initial mixing. Water or irrigate after application to improve penetration of the fungicide into the soil.

### Application

Use only at recommended rates, as overdosage may be harmful to sensitive plants. A test for compatibility with other chemicals is advisable before use.

## FUNGICIDE RESISTANCE WARNING

### GROUP 14 FUNGICIDE

For fungicide resistance management, Terrazole 350 WP Soil Fungicide is a Group 14 fungicide. Some naturally occurring individual fungi resistant to Terrazole 350 WP Soil Fungicide and other Group 14 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal populations if these fungicides are used repeatedly. These resistant fungi will not be controlled by Terrazole 350 WP Soil Fungicide or other Group 14 fungicides, thus resulting in a reduction in efficacy and possible yield loss.

Since the occurrence of resistant fungi is difficult to detect prior to use, Arysta LifeScience Australia Pty Ltd accepts no liability for any losses that may result from the failure of Terrazole 350 WP Soil Fungicide to control resistant fungi.

### PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT plant food crops in rotation on treated soils.

### PROTECTION OF LIVESTOCK

DO NOT allow stock to graze in any treated area. DO NOT feed turf/lawn clippings from any treated area to poultry or livestock.

### PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Dangerous to fish and other aquatic organisms. DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

## STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

## SAFETY DIRECTIONS

Avoid contact with eyes and skin. Do not inhale dust or spray mist. Wash hands after use.

## FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

## MATERIAL SAFETY DATA SHEET

For additional information, refer to the Material Safety Data Sheet.

## CONDITIONS OF SALE

Arysta LifeScience Australia Pty Ltd will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply or use of these goods other than responsibility of the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Arysta LifeScience Australia Pty Ltd is limited to the replacement of the goods or (at the option of Arysta LifeScience Australia Pty Ltd) the refund of the price paid and is conditional upon a claim being made in writing and where possible sufficient part of the goods to enable proper examination being returned to Arysta LifeScience Australia Pty Ltd within thirty days of delivery.

®Terrazole is a registered trademark of MacDermid Agricultural Solutions Inc.

BN  
DOM

APVMA Approval No. 53332/10p/1206



09/15/B