

POISON

KEEP OUT OF REACH OF CHILDREN

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

*O-Mat*TM 290 SL

INSECTICIDE

**ACTIVE CONSTITUENT: 290 g/L OMETHOATE
(AN ANTICHOLINESTERASE COMPOUND)**

GROUP 1B INSECTICIDE

For control of redlegged earth mite, blue oat mite and lucerne flea on pastures, cereals, oilseed and legume crops, and bluegreen aphid and cowpea aphid on pasture legumes, lucerne, faba beans and vetch as specified in the Directions for Use Table.

**IMPORTANT: PLEASE READ THE ATTACHED BOOKLET
BEFORE USING THIS PRODUCT**

Contents: 10 L

Arysta LifeScience Australia Pty Ltd
Level 3, 70 Hindmarsh Square, Adelaide SA 5000
Tel: (08) 8112 0900 Fax: (08) 8112 0999



 **Arysta**
LifeScience



DIRECTIONS FOR USE

CROP	PEST	RATE	CRITICAL COMMENTS
Pastures, Cereals, Oilseed and Legume crops	Redlegged Earth Mite (Not Qld)	100 mL/ha	AUTUMN/WINTER: Spray 2–5 weeks after opening rains when mites have hatched to prevent seedling damage or loss. O-Mat is rain fast within 1 hour of spraying. Residual control is less on 1–2 true leaf plants. O-Mat should be mixed with minimum tillage or direct drill sprays to kill existing mite populations. SPRING: Spray mite infested pastures and legumes to prevent damage, yield loss and to reduce infestation pressure in the following autumn. O-Mat may be added to pasture herbicides.
	Blue Oat Mite (Qld, NSW only)		
	Lucerne Flea (Not Qld)		Spray on appearance and before damage occurs. If second hatching occurs a follow up spray may be required.
	Pasture Mite (Bryobia Mite) (NSW, WA only)	120 mL/ha	Spray on appearance and before damage occurs.
Pasture legumes, Lucerne, Faba bean, Vetch	Bluegreen Aphid	100 or 200 mL/ha	Spray when aphids start to build up. Use the higher rate when faster knockdown is required.
	Cowpea Aphid	200 mL/ha	Spray when aphids start to build up.
Pasture	Spotted Clover Aphid	100 mL/ha	Apply at first sign of aphid activity.
Barrier spraying	Redlegged earth mite (NSW, WA only)	300 mL/ha	To prevent invasion, spray bare earth outside crop as well as weeds along the fenceline using an offset jet. Note: Rain will reduce the efficacy of the bare earth treatment but leave an effective barrier on the weeds.
Poppy	Redlegged Earth Mite (Tas only)	100 mL/ha	Spray on appearance and before damage occurs.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

WITHHOLDING PERIOD: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 1 DAY AFTER APPLICATION

GENERAL INSTRUCTIONS**Insecticide Resistance Warning**

GROUP 1B	INSECTICIDE
-----------------	--------------------

For insecticide resistance management, O-Mat is a Group **1B** insecticide. Some naturally occurring insect biotypes resistant to O-Mat and other Group **1B** insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if O-Mat and other Group **1B** insecticides are used repeatedly. The effectiveness of O-Mat on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, Arysta LifeScience Corporation accepts no liability for any losses that may result from the failure of O-Mat to control resistant insects. O-Mat may be subject to specific resistance management strategies. For further information contact your local supplier, Arysta LifeScience representative or local agricultural department agronomist.

Mixing

Mix the required quantity of O-Mat in sufficient water to give even coverage. Boom sprayers and aircraft are satisfactory for spraying O-Mat.

Compatibility

Compatible with most commonly used herbicides from direct drilling and minimum tillage to pasture topping. Ensure that O-Mat is thoroughly mixed in the spray tank before adding a herbicide. DO NOT mix concentrates together but add each to the spray tank separately. As formulations of other manufacturers' products are beyond the control of Arysta LifeScience Corporation, all mixtures should be tested prior to mixing commercial quantities. As changes in climatic conditions can alter the sensitivity of plants to mixtures of sprays, Arysta LifeScience cannot be responsible for the behaviour of such mixtures.

PROTECTION OF LIVESTOCK

Dangerous to bees. DO NOT spray any plants in flower while bees are foraging.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse container before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, State or Territory government regulations. DO NOT burn empty container or product.

SAFETY DIRECTIONS

Product and spray are poisonous if absorbed by skin contact, inhaled or swallowed. Will irritate the eyes. Repeated exposure may cause allergic disorders. Repeated minor exposure may have a cumulative poisoning effect. Avoid contact with eyes and skin. DO NOT inhale spray mist. When preparing spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC

gloves and face shield and impervious footwear. If clothing becomes contaminated with product remove clothing immediately. If product on skin, immediately wash area with soap and water. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands arms and face thoroughly with soap and water. After each day's use wash gloves, face shield and contaminated clothing.

FIRST AID

If swallowed, splashed on skin or in eyes, or inhaled, contact a Poisons Information Centre (Phone Australia 131126) or a doctor at once. Remove any contaminated clothing and wash skin thoroughly.

If swallowed, activated charcoal may be advised. Give atropine if instructed.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet, which can be obtained from Arysta LifeScience Australia Pty Ltd.

EXCLUSION OF LIABILITY

Arysta LifeScience Australia Pty Ltd has no control over storage, handling and manner of use of this product. Where this material is not stored, handled or used correctly and in accordance with the directions, no express or implied representations or warranties concerning this product (other than non-excludable statutory warranties) will apply. Arysta LifeScience Pty Ltd accepts no responsibility for any loss arising from incorrect storage, handling or use.

™ Arysta LifeScience Australia Pty Ltd

APVMA Approval No. 82095/104990