

**POISON**  
**KEEP OUT OF REACH OF CHILDREN**  
**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

*Arysta LifeScience*

# **LVE MCPA 570**

**HERBICIDE**

**ACTIVE CONSTITUENT: 570g/L MCPA,  
present as the 2-ETHYLHEXYL ESTER**

**GROUP I HERBICIDE**

**For selective control of certain weeds in agricultural  
crops as listed in the Directions for Use.**

**IMPORTANT: READ THE ATTACHED LEAFLET  
BEFORE USING THIS PRODUCT.**

**Contents: 20 L**

Arysta LifeScience Australia Pty Ltd  
Level 7, 435 King William Street Adelaide SA 5000  
Tel: (08) 8112 0900 Fax: (08) 8112 0999

08/15/F



File name	LVE MCPA 570 Label 20L.pdf
Dimensions	Size 170mm H x 170mm W
Scale	Prints at 100% A4
Date	13/08/2015

## DIRECTIONS FOR USE:

Situation & Crop	Weeds	Rate/ha	Critical Comments
Wheat (up to early flag leaf (Z37) only)	Refer Weed Table	440 mL – 1.84 L	Use the lower rate (440 mL/ha) when crop is at 3–5 leaf stage (Z13-Z15). Apply higher rates from 5 leaf stage through to flag leaf just visible (Z15-Z37). <b>Kulin:</b> Use only from Z15-Z37 (WA only).
Oats, Barley, Triticale and Cereal Rye (up to early flag leaf (Z37) only)		440 mL – 1.84 L	Use lower rate (440 mL/ha) when crop is at 3–5 leaf stage (Z13-Z15). Apply higher rates from 5 leaf stage through to flag leaf just visible (Z15-Z37). Barley: Use only from 5 leaf stage to flag leaf just visible (Z15-Z37).
Grass Pastures and Grass Seed Crops		440 mL – 1.84 L	Apply to established grass based pastures. Damage may occur to Legumes if present. See Legume Tolerance under Protection of Crops. <b>NOTE:</b> Capeweed, Paterson's Curse (Salvation Jane) and Variegated Thistle may be poisonous to stock after spraying.
<b>CEREALS</b> Wheat, barley, oats, triticale, cereal rye	<b>Wild radish</b> Stage of weed growth	ARYSTA LIFESCIENCE LVE MCPA 570 HERBICIDE 175 mL plus LV MCPA/ Diflufenican Selective Herbicide 350 mL	*Reduced efficacy (suppression only) may be achieved on wild radish larger than 8 leaf or greater than 180 mm in diameter. DO NOT use this tank-mix if cereals are undersown with lucerne or annual medics. <b>Crop Stage</b> LV MCPA/Diflufenican Selective Herbicide 350 mL + ARYSTA LIFESCIENCE LVE MCPA 570 HERBICIDE 175 mL: Apply from 3 leaf to fully tillered (Z13–30).
	Up to the 4 leaf stage and not more than 120 mm in diameter	ARYSTA LIFESCIENCE LVE MCPA 570 HERBICIDE 175 mL plus LV MCPA/ Diflufenican Selective Herbicide 500 mL	LV MCPA/Diflufenican Selective Herbicide 500 mL + ARYSTA LIFESCIENCE LVE MCPA 570 HERBICIDE 175 mL: Apply from 3 leaf to fully tillered (Z13–30).
	Up to the 6 leaf stage and not more than 150 mm in diameter	ARYSTA LIFESCIENCE LVE MCPA 570 HERBICIDE 175 mL plus LV MCPA/ Diflufenican Selective Herbicide 500 mL	LV MCPA/Diflufenican Selective Herbicide 500 mL + ARYSTA LIFESCIENCE LVE MCPA 570 HERBICIDE 350 mL: Apply from 5 leaf stage to fully tillered (Z16–30).
	Up to the 8 leaf stage and not more than 180 mm in diameter*	ARYSTA LIFESCIENCE LVE MCPA 570 HERBICIDE 350 mL plus LV MCPA/ Diflufenican Selective Herbicide 500 mL	Optimum results are achieved when sprayed at 3–5 leaf crop stage (generally 4–8 weeks post-sowing). <b>WA only:</b> DO NOT apply to Barley or Kulin Wheat before the 5 leaf stage (Z15). <b>Warning:</b> LV MCPA/Diflufenican Selective Herbicide may cause transient crop yellowing of cereals. Some varieties of oats have not been tested. (Refer to "Crop Tolerance" section of General Instructions on the LV MCPA/Diflufenican Selective Herbicide label).

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**WITHHOLDING PERIOD:**

**HARVEST: NOT REQUIRED WHEN USED AS DIRECTED.**

**GRAZING: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION.**

**WEED TABLE:** Specific Weed Control Recommendations.

Where weeds are to be sprayed in a crop or pasture, use only the rates given for the particular crop or situation indicated under the Directions of Use table.

In most cases this will give control, however some hard to kill weeds or those in advanced stages of growth may only be suppressed.

The rates listed in the weeds table below are for use where weeds only are present and crop or pasture is not involved.

Use at these rates in a crop or pasture other than for spot spraying may cause damage.

Weeds	Rate/ha	Critical Comments
<b>Bathurst Burr</b> ( <i>Xanthium spinosum</i> )	1.4 L	Moderately susceptible. Spray young seedlings.
<b>Capeweed</b> ( <i>Arctotheca calendula</i> )	1.49 L	Apply up to 2–4 leaf stage only when plants are no larger than 5 cm in diameter.
<b>Charlock</b> ( <i>Sinapis arvensis</i> )	400 mL–1.4 L	Young rosettes very susceptible.
<b>Dandelion</b> ( <i>Taraxacum officinale</i> )	1.4 L	Apply in early rosette stage.
<b>Fat Hen</b> ( <i>Chenopodium album</i> )	880 mL–1.4 L	Tolerance increases with age. Use lower rate only on seedlings.
<b>Fumitory</b> ( <i>Fumaria</i> )	965 mL	Moderately susceptible. Spray up to first true leaf stage. (Not Tas).
<b>Hedge Mustards</b> ( <i>Sisymbrium</i> spp)	490–880 mL	Apply at rosette stage.
<b>Lincoln Weed</b> ( <i>Diploaxis tenuifolia</i> )	615 mL	Apply up to rosette stage only.
<b>London Rocket</b> ( <i>Sisymbrium irio</i> )	965 mL	Spray at young stage.
<b>Noogoora Burr</b> ( <i>Xanthium occidentale</i> )	1.4 L	Tolerance increases with age. Spray young seedlings only.
<b>Paterson's Curse</b> ( <i>Echium plantagineum</i> )	1.49 L	Apply at 2 to 4 leaf stage only.
<b>Saffron Thistle</b> ( <i>Carthamus lanatus</i> )	965 mL–1.66 L	Spray young rosettes before leaves become spiny.
<b>Scotch (Cotton) Thistle</b> ( <i>Onopordum acanthium</i> )	615 mL–1.49 L	Spray young rosettes.
<b>Skeleton Weed</b> ( <i>Chondrilla juncea</i> )	965 mL–1.4 L	Moderately susceptible. Spray rosettes before aerial growth commences.
<b>Slender Thistle</b> ( <i>Carduus tenuiflorus</i> )	1.84 L	Moderately susceptible. Spray young rosettes.
<b>Spear Thistle</b> ( <i>Cirsium vulgare</i> )	1.31–1.84 L	Moderately susceptible. Spray young rosettes.
<b>Stinkwort</b> ( <i>Dittrichia graveolens</i> )	1.4 L	Spray very young before plants 15 cm high.
<b>Turnip Weed</b> ( <i>Rapistrum rugosum</i> )	615–965 mL	Apply up to rosette stage. Apply lower rates to seedlings.
<b>Volunteer Canola</b> ( <i>Brassica napus</i> )	1.31 L	Apply to seedlings only.
<b>Variiegated Thistle</b> ( <i>Silybum marianum</i> )	735–965 mL	Spray at pre-cabbage stage.
<b>Volunteer Safflower</b> ( <i>Carthamus tinctorius</i> )	880 mL	Apply to seedlings only.
<b>Volunteer Sunflower</b> ( <i>Helianthus annuus</i> )	1.31 L	Apply to seedlings only.
<b>Wild Radish</b> ( <i>Raphanus raphanistrum</i> )	965–1.4 L	Apply up to 2 to 3 leaf stage only.
<b>Wild Sage</b> ( <i>Salvia verbenaca</i> )	965 mL	Spray young seedlings.
<b>Wild Turnip</b> ( <i>Brassica</i> spp)	440 mL–1.31 L	Spray young rosettes. Lower rates for seedlings.

## **GENERAL INSTRUCTIONS**

Before opening, carefully read Directions for Use, Precautionary Statements, Safety Directions and First Aid Instructions. Weeds should be sprayed while actively growing and at their most susceptible stage. As a general guide, annuals should be sprayed when young and perennials just prior to appearance of bud stage. Extremes of cold or drought are unfavourable. The best conditions are when soil is moist, weather fine and rain unlikely within 6 hours following application. DO NOT spray weeds outside the stages indicated in “Critical Comments” as damage, loss of yield or inadequate weed control may result. DO NOT spray in high winds.

## **APPLICATION/MIXING INFORMATION**

This product mixes readily with water. Half fill the spray tank with clean water and add the required amount of product. Agitate thoroughly before adding remainder of water.

BOOM SPRAYING—Use 30–120 L/ha of water.

AERIAL SPRAYING—Use 10–90 L/ha of water.

## **EQUIPMENT MAINTENANCE AND USAGE**

Equipment that has been used for this chemical should not be used for the application of other materials to sensitive plants unless it has been well washed out with Spraymate Tank & Equipment Cleaner followed by several clear water rinses.

## **COMPATIBILITY/TANK MIXES**

This product can be mixed with ARYSTA LIFESCIENCE LVE MCPA 570 HERBICIDE, Kamba®, Bromicide®, Flowable Diuron, Flowable Terbutryn, Glean\*/Siege\*, Arysta LifeScience Glyphosate, Roundup ^, Grasp\*, Diclofop-Methyl 375 Selective Herbicide, Triasulfuron 750 WG Herbicide, LV MCPA/Diflufenican Selective Herbicide, Nugrex/Tigrex\*, Lontrel\*/Archer™, Propon, Puma\*. When tank mixing add the required amount of compatible product when the spray tank is nearly full and while continuously agitating. If left overnight, agitate thoroughly before spraying. When mixing with Diclofop-Methyl 375 EC Selective Herbicide add an approved Wetter 1000 surfactant and do not exceed 700 mL of ARYSTA LIFESCIENCE LVE MCPA 570 HERBICIDE per hectare. For consistent results, use a minimum of 80 L water per ha by ground and 30 L by air, and if wild oats are present, use 2 L Diclofop-Methyl 375 EC Selective Herbicide per hectare.

**GROUP****I****HERBICIDE****RESISTANT WEEDS WARNING**

ARYSTA LIFESCIENCE LVE MCPA 570 HERBICIDE is a member of the Phenoxy group of herbicides. ARYSTA LIFESCIENCE LVE MCPA 570 HERBICIDE has the Disruptors of plant cell growth mode of action. For weed resistance management ARYSTA LIFESCIENCE LVE MCPA 570 HERBICIDE is a Group I herbicide. Some naturally-occurring weed biotypes resistant to ARYSTA LIFESCIENCE LVE MCPA 570 HERBICIDE and other Group I herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by ARYSTA LIFESCIENCE LVE MCPA 570 HERBICIDE or other Group I herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Arysta LifeScience Australia Pty Ltd accepts no liability for any losses that may result from the failure of ARYSTA LIFESCIENCE LVE MCPA 570 HERBICIDE to control resistant weeds.

**PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS**

DO NOT apply under weather conditions or from spraying equipment that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures. Avoid spray drift and vapour movement onto susceptible crops such as cotton, vines, tomatoes, vegetables, fruit trees, lucerne, legume crops and pastures, oilseed crops or other susceptible crops or trees (e.g. kurrajongs, belahs, eucalypts).

**Legume Tolerance:** The tolerance of pasture legumes to this product can vary with species, variety, growing conditions, stage of growth and companion crop cover. While MCPA is preferable to 2,4-D in most situations, this product will cause greater damage than MCPA amine formulations. Lucerne and medics should not be sprayed under any circumstances with this product as severe damage will result.

**PROTECTION OF WILDLIFE, FISH, CRUSTACEA AND ENVIRONMENT**

DO NOT contaminate streams, rivers or waterways with the chemical or used container.

**STORAGE AND DISPOSAL** Store in the closed, original container in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemical on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. Do not burn empty containers or product.

**Refillable containers (1000 L only):**

Store in the closed, original container in a cool well ventilated area. DO NOT store for prolonged periods in direct sunlight. Empty contents fully into application equipment. Close all valves and return to point of sale for refill or storage.

**Envirodrum 110 L Mini Bulk Returnable Container:**

Store the original sealed Envirodrum™ in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight. DO NOT tamper with the Micro Matic valve or the security seal. DO NOT contaminate the Envirodrum with water or any foreign matter. After each use of the product, please ensure that the Micro Matic coupler, delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the Envirodrum have been used, please return the empty Envirodrum to the point of purchase. The Envirodrum remains the property of Arysta LifeScience Australia Pty Ltd.

**SAFETY DIRECTIONS**

Harmful if swallowed. Attacks eyes. Will irritate the skin. When preparing spray and using the prepared spray wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and goggles. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms, and face thoroughly with soap and water. After each day's use wash gloves, goggles and contaminated clothing.

**FIRST AID**

If poisoning occurs, contact a Doctor or Poisons Information Centre (Phone Australia 13 11 26).

**MATERIAL SAFETY DATA SHEET**

For further information refer to the Material Safety Data Sheet (MSDS).

**CONDITIONS OF SALE**

"Any provisions or rights under the Trade Practices Act 1974 or relevant state legislation which cannot be excluded by those statutes or by law are not intended to be excluded by these conditions of sale. Subject to the foregoing, all warranties, conditions, rights and remedies, expressed or implied under common law, statute or otherwise, in relation to the sale, supply, use or application of this product, are excluded. Arysta LifeScience Australia Pty Ltd and/or its affiliates ("Arysta LifeScience") shall not accept any liability whatsoever (including consequential loss), or howsoever arising (including negligence) for any damage, injury or death connected with the sale, supply, use or application of this product except for liability which cannot be excluded by statute."

®, ™, ^, \* **Registered trademarks**

<b>UN No. 3082</b>	<b>ENVIRONMENTALLY HAZARDOUS SUBSTANCE, LIQUID, N.O.S. (CONTAINS MCPA-2-Ethylhexyl Ester) MARINE POLLUTANT</b>
In a Transport Emergency Dial <b>000</b> Police or Fire Brigade	SPECIALIST ADVICE IN EMERGENCY ONLY <b>1800 033 111</b> ALL HOURS – AUSTRALIA WIDE
<b>P.G. III</b>	<b>HAZCHEM 3[Z]</b>

