

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Thiraflo[®]
FLOWABLE FUNGICIDE



ACTIVE CONSTITUENT: 600 g/L THIRAM

GROUP M3 FUNGICIDE

For use as a seed treatment to control various fungal diseases of seeds and as a drench to control fungal diseases of turf as specified in the Directions for Use table.

IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USE

Contents: 20 L

Arysta LifeScience Australia Pty Ltd
Level 7, 435 King William Street Adelaide SA 5000
Tel: (08) 8112 0900 Fax: (08) 8112 0999

08/15/F



POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE
OPENING OR USING

Thiraflo
FLOWABLE FUNGICIDE



ACTIVE CONSTITUENT: 600 g/L THIRAM

GROUP M3 FUNGICIDE

For use as a seed treatment to control various fungal diseases of seeds and as a drench to control fungal diseases of turf as specified in the Directions for Use table.

IMPORTANT: READ THIS LEAFLET BEFORE USE

Arysta LifeScience Australia Pty Ltd
Level 7, 435 King William Street
Adelaide SA 5000
Tel: (08) 8112 0900 Fax: (08) 8112 0999



DIRECTIONS FOR USE:

RESTRAINTS:

DO NOT mix this product in a slurry with rhizobial inoculants. Only treat good quality sound seed as treatment of mechanically damaged or poor quality seed may result in reduced germination and/or reduced seedling vigour and emergence.

| Situation | Disease | Rate | Critical Comments |
|--|---|--|--|
| Turf | Brown patch (<i>Rhizoctonia solani</i>) Helminthosporium disease (<i>Helminthosporium</i> spp.) Fusarium patch (<i>Fusarium</i> spp.) | 160 mL in 20 L of water per 100 m ² | Apply at first sign of disease and repeat at 7–14 day intervals, particularly in humid weather. |
| | Damping off (<i>Pythium</i> spp.) | 200 mL in 60 L of water per 100 m ² | Apply as a drench to the seedbed. Repeat every 5–7 days during disease susceptible periods. |
| Sorghum | Seedling decay complex (<i>Pythium</i> and <i>Rhizoctonia</i> spp.) | 2.5–3.5 L per tonne seed | THIRAFLO must be applied at the correct rate to achieve uniform coverage of all seeds. THIRAFLO can be applied to the seed undiluted or mixed with water to achieve optimum coverage of grain with the seed treatment. Use the higher rate when disease pressure is expected to be severe. If water is required, make a slurry by mixing 2.7 L of THIRAFLO with 2.3 L of water. Apply this mixture per tonne of seed. Apply only to good quality graded seed. Move the seed up the grain auger or apply in a suitable mixer. The mixing of seed after application is essential to obtain good seed coverage. The total volume of slurry applied per tonne should be minimised to approximately 5 L. In the case of ascochyta and botrytis, THIRAFLO will reduce the incidence of seed-borne infection, but follow up foliar fungicide sprays registered for this purpose will be required to control field infection. |
| Chickpea seed | Seed-borne botrytis (<i>Botrytis</i> spp.) Seed-borne ascochyta blight (<i>Ascochyta rabiei</i>) | 2.0 L per tonne seed | |
| Lupin seed | Seed-borne anthracnose (<i>Colletotrichum</i> spp.) | 1.7–2.0 L per tonne seed | |
| Sorghum and sunflower seed – for export only | To meet export requirements | Up to 4 L per tonne of seed | Treated seed is for export only. Use of product must be in accordance with importing countries requirements. Be aware of any importing countries residue tolerance limits and do not apply product at a rate that will exceed the tolerance limits of the importing country. Treated seed that is not exported must be disposed of in a manner that does not result in planting or consumption by humans or animals. |

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD:

NOT REQUIRED WHEN USED AS DIRECTED

GENERAL INSTRUCTIONS

THIRAFLO FLOWABLE FUNGICIDE is a suspension concentrate (flowable) fungicide for use as a seed treatment to control various fungal diseases of seeds and as a drench to control fungal diseases of turf.

INOCULATION OF SEED

This product does not adversely affect germination if used as directed; however, when rhizobial inoculants are also applied to seed, the viability of the inoculum may be reduced. Inoculum can be applied to the seed in a separate operation after seed treatment. Apply the inoculum on the day of sowing after the fungicide treatment has dried on the seed. Sow into a moist seedbed as soon as possible after treatment. If seed is sown into a dry seedbed, then fungicidal seed treatment can cause a reduction in nodulation. Alternatively, inoculum can be sprayed into the furrow at seeding rather than combined on seed treated with this product. This method of applying inoculum is not recommended in Western Australia.

MIXING AND APPLICATION

Turf should be in good nutritional condition prior to treatment to prevent damage to sensitive turf species (eg fescues, bluegrass and turf type ryegrass).

THIRAFLO FLOWABLE FUNGICIDE may be applied to seed undiluted or may be mixed with water if required to achieve optimum coverage of seed. This requirement will depend on the equipment in use and the rates to be applied. Use only equipment designed to apply suspension concentrate or slurry seed treatments to seed. As the product can dry out on exposed internal container surfaces under hot summer conditions resulting in the formation of flakes, it should be filtered before use. Do not let liquid dry out in application equipment as flakes can block filters and limit the flow to spray nozzles. Equipment used for applying this product can be cleaned out with water. Shake the container before adding water to the product. After adding the required amount of water to the product, shake again. Where the diluted product has been permitted to stand for an extended period, then the mixture should again be agitated well.

Handle seed carefully after application and avoid excess wetting to prevent damage to the seed coat.

SOWING

Check seed flow through the drill after treatment as treated seed can flow more slowly.

STORAGE OF TREATED SEED

If seed is not used immediately after treatment, it should be stored in a dry, well-ventilated place. Although THIRAFLO FLOWABLE FUNGICIDE has no effect on the viability of treated seed, subsequent germination can be adversely affected by poor storage conditions such as high moisture combined with high temperatures. No liability can therefore be accepted for performance of stored treated seed. Treated seed should not be held over from one sowing season for sowing in subsequent seasons.

FUNGICIDE RESISTANCE WARNING

GROUP M3 FUNGICIDE

For fungicide resistance management, THIRAFLO FLOWABLE FUNGICIDE is a Group M3 fungicide. Some naturally occurring individual fungi resistant to THIRAFLO FLOWABLE FUNGICIDE and other Group M3 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal populations if these fungicides are used repeatedly. These resistant fungi will not be controlled by THIRAFLO FLOWABLE FUNGICIDE or other Group M3 fungicides, thus resulting in a reduction in efficacy and possible yield loss.

Since the occurrence of resistant fungi is difficult to detect prior to use, Arysta LifeScience Australia Pty Ltd accepts no liability for any losses that may result from the failure of THIRAFLO FLOWABLE FUNGICIDE to control resistant fungi.

PROTECTION OF LIVESTOCK

Seed treated with THIRAFLO FLOWABLE FUNGICIDE must not be used for animal or human consumption. It is important that any treated seed not used for sowing, is destroyed. Bags that have held treated seed should not be used for any other purpose. DO NOT allow seed treated with THIRAFLO FLOWABLE FUNGICIDE to contaminate grain intended for animal or human consumption. When treated seed is stored, it should be left apart from untreated grain and the bags (or other containers) should be clearly marked to indicate the contents have been treated with THIRAFLO FLOWABLE FUNGICIDE.

DO NOT feed or otherwise expose treated seed to domestic animals or birds.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT feed or otherwise expose treated seed to wild animals and birds. Dangerous to fish. DO NOT contaminate streams, rivers or waterways and drains with this product or used containers/bags that have held treated seed.

STORAGE AND DISPOSAL

Store in the closed, original container in a well-ventilated area, as cool as possible. Do not store for prolonged periods in direct sunlight.

This container can be recycled if it is clean, dry, free of visible residues and has the drumMUSTER logo visible. Triple-rinse container for disposal. Dispose of rinsate by adding it to the spray tank. Do not dispose of undiluted chemical on site. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any drumMUSTER collection or similar container management program site. The cap should not be replaced, but may be taken separately.

SAFETY DIRECTIONS

May irritate the eyes, nose, throat and skin. Repeated exposure may cause allergic disorders. Avoid contact with eyes and skin. Do not inhale spray mist. When opening the container, preparing the product and using the prepared product wear cotton overalls buttoned to the neck and wrist, elbow length rubber gloves, face shield or goggles. If product on skin, immediately wash area with soap and water. If product in eyes wash it out immediately with water. Wash hands after use. After each days use, wash gloves, face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre on 13 11 26 (Australia). Avoid giving alcohol.

MATERIAL SAFETY DATA SHEET

For additional information, refer to the Material Safety Data Sheet.

CONDITIONS OF SALE

Arysta LifeScience Australia Pty Ltd will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply or use of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Arysta LifeScience Australia Pty Ltd is limited to the replacement of the goods or (at the option of Arysta LifeScience Australia Pty Ltd) the refund of the price paid and is conditional upon a claim being made in writing and where possible sufficient part of the goods to enable proper examination being returned to Arysta LifeScience Australia Pty Ltd within thirty days of delivery.

® Thiraflo is a registered trademark of Arysta LifeScience Australia Pty Ltd.

STORAGE AND DISPOSAL

Store in the closed, original container in a well-ventilated area, as cool as possible. Do not store for prolonged periods in direct sunlight.

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemical on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

May irritate the eyes, nose, throat and skin. Repeated exposure may cause allergic disorders. Avoid contact with eyes and skin. Do not inhale spray mist. When opening the container, preparing the product and using the prepared product wear cotton overalls buttoned to the neck and wrist, elbow length rubber gloves, face shield or goggles. If product on skin, immediately wash area with soap and water. If product in eyes wash it out immediately with water. Wash hands after use. After each days use, wash gloves, face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre on 131 126 (Australia).
Avoid giving alcohol.

MATERIAL SAFETY DATA SHEET



For additional information, refer to the Material Safety Data Sheet.

CONDITIONS OF SALE

Arysta LifeScience Australia Pty Ltd will not accept any responsibility whatsoever and howsoever arising and whether for consequential loss or otherwise in connection with the supply or use of these goods other than responsibility for the merchantable quality of the goods and such responsibilities mandatorily imposed by Statutes applicable to the sale or supply of these goods. To the extent allowed by such Statutes the liability of Arysta LifeScience Australia Pty Ltd is limited to the replacement of the goods or (at the option of Arysta LifeScience Australia Pty Ltd) the refund of the price paid and is conditional upon a claim being made in writing and where possible sufficient part of the goods to enable proper examination being returned to Arysta LifeScience Australia Pty Ltd within thirty days of delivery.

* Thiram is a registered trademark of Arysta LifeScience Australia Pty Ltd.

| | |
|--|---|
| UN No. 3082 | ENVIRONMENTALLY HAZARDOUS SUBSTANCE, LIQUID, N.O.S. (CONTAINS THIRAM) MARINE POLLUTANT |
| In an Emergency Dial 000 Police or Fire Brigade | SPECIALIST ADVICE IN EMERGENCY ONLY 1800 033 111 ALL HOURS – AUSTRALIA WIDE |
| P.G. III | HAZCHEM 2X |



Batch Number:

Date of Manufacture:

APVMA Approval No. 58072/20L/1109

9 322524 067095