

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Arysta LifeScience

Tebcon

430 SC FUNGICIDE

ACTIVE CONSTITUENT: 430 g/L TEBUCONAZOLE

GROUP 3 FUNGICIDE

For the control of leaf spot and leaf speckle on bananas; rust, leaf spot and net blotch of peanuts; foliar diseases on cereal crops; and other diseases on beans, peas, onions, pawpaw, pyrethrum and ryegrass and fescue seed crops as specified in the Directions for Use Table.

IMPORTANT: PLEASE READ THE ATTACHED LEAFLET BEFORE USING THIS PRODUCT

Contents: 20 L

Arysta LifeScience Australia Pty Ltd
Level 3, 70 Hindmarsh Square, Adelaide SA 5000
Tel: (08) 8112 0900 Fax: (08) 8112 0999



 **Arysta**
LifeScience

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Arysta LifeScience

Tebcon

430 SC FUNGICIDE

ACTIVE CONSTITUENT:
430 g/L TEBUCONAZOLE

GROUP 3 FUNGICIDE

For the control of leaf spot and leaf speckle on bananas; rust, leaf spot and net blotch of peanuts; foliar diseases on cereal crops; and other diseases on beans, peas, onions, pawpaw, pyrethrum and ryegrass and fescue seed crops as specified in the Directions for Use Table.

IMPORTANT: READ THIS LEAFLET BEFORE USE

Arysta LifeScience Australia Pty Ltd
Level 3, 70 Hindmarsh Square
Adelaide SA 5000
Tel: (08) 8112 0900 Fax: (08) 8112 0999



DIRECTIONS FOR USE

CROP	DISEASE	APPLICATION	RATE mL/ha	WHP	CRITICAL COMMENTS
Bananas (Qld, NSW, WA, NT only)	Leaf spot (Yellow sigatoka) Leaf speckle Black Sigatoka	Ground Application Apply by misting machine or airstream sprayer in a convenient volume of water Aerial application Apply at least 20 L of spray mixture per hectare	230 Add 3–6 L water miscible oil per ha	H 1 day	Maintain good de-leafing practices to reduce disease inoculum. Very old leaves and leaves with advanced lesions should be removed, or infected parts of the leaf removed, prior to the application of ARYSTA LIFESCENCE TEBCON 430 SC FUNGICIDE. Tropical Areas (e.g. North Qld, NT, Ord River): Apply a regular schedule of protectant sprays. When conditions favour disease, apply a maximum of 2 consecutive ARYSTA LIFESCENCE TEBCON 430 SC FUNGICIDE sprays at 14 day intervals. Do not apply more than 6 ARYSTA LIFESCENCE TEBCON 430 SC FUNGICIDE sprays in any 12 month period. Do not apply any ARYSTA LIFESCENCE TEBCON 430 SC FUNGICIDE sprays in the months of June, July, August and September. Sub-Tropical Areas (e.g. South Qld, NSW): Commence spraying with ARYSTA LIFESCENCE TEBCON 430 SC FUNGICIDE at the onset of warm and humid/wet weather, normally December. Repeat at 21 to 28 day intervals using a maximum of 2 consecutive ARYSTA LIFESCENCE TEBCON 430 SC FUNGICIDE sprays. Do not apply more than 5 ARYSTA LIFESCENCE TEBCON 430 SC FUNGICIDE sprays in any 12 month period. ARYSTA LIFESCENCE TEBCON 430 SC FUNGICIDE is approved for use in banana plantations inter-planted with avocados.
Peanuts (South Qld and NSW only)	Early Leaf spot Late leaf spot Rust Net Blotch	Low Disease Ground Application Apply in at least 100 L of water per ha Aerial application Apply at least 30 L of spray mixture per hectare High disease OR Wet weather High disease AND Wet weather	175 + Agridex 1 L/ha 290 + Agridex 1 L/ha 440 + Agridex 1 L/ha	H 3 weeks G 3 weeks	Regularly check high-risk areas in the crop for disease (e.g. lower leaves, shaded plants). When leaf spot or net blotch or rust can easily be found, then either: • Spray immediately after last disease conducive weather (e.g. rain or heavy dews), or • Spray before the next conducive weather. Repeat after 14 days if conditions remain favourable to disease development. If not, repeat just before or just after the next disease conducive weather. For resistance strategy, see General Instructions.
Peanuts (North Qld, WA, NT only)	Early Leaf spot Late Leaf spot Rust	Low Disease Moderate disease Severe disease	230 + Agridex 1 L/ha 290 + Agridex 1 L/ha 440 + Agridex 1 L/ha		Begin spraying at 3–4 weeks after planting. If band spraying, apply half the appropriate rate as a 45cm band directly over the row. Sprays should not be banded after 6 weeks from planting. Repeat applications at 14-day intervals. If prolonged wet weather or heavy rains occur, shorten spray interval to 10–12 days. For resistance strategy, see General Instructions.
	Net blotch	Low Disease Moderate–severe disease	290 + Agridex 1 L/ha 440 + Agridex 1 L/ha		Apply at 14 day intervals. If prolonged cool moist weather occurs, shorten spray interval to 10–12 days. For resistance strategy, see General Instructions.

CROP	DISEASE	APPLICATION	RATE mL/ha	WHP	CRITICAL COMMENTS
Green Beans	Rust	Can be applied by aircraft or ground rig	350 + Agridex 1 L/ha	H 3 days G 3 weeks	Spray when rust infection begins or at budding, whichever is the earlier. Repeat application 10–14 days later. A third application may be necessary when infection occurs early or disease pressure is high.
Peas	Powdery Mildew	Ground Application Apply in at least 50 L of water per ha Aerial application Apply in at least 10 L of spray mixture per hectare	145 290		Apply at flowering or at the first sign of disease, whichever occurs first. A second spray 14 days later may be necessary under some conditions.
Wheat	Leaf rust Stripe rust Septoria nodorum blotch Yellow leaf spot Septoria tritici blotch		145 or 290 290	H 5 weeks G 14 days	Use the higher rate when longer disease control is required. Stripe rust: See spray timings under General Instructions. Other diseases: Apply from full flag leaf emergence to early head emergence. The addition of mineral crop oil (e.g. DC Trate or equivalent) at 1% may improve performance of ARYSTA LIFESCENCE TEBCON 430 SC FUNGICIDE on wheat, oats and barley.
Oats	Crown rust		145 or 290		Apply at late tillering to early jointing.
Barley	Scald		145		Apply when 5% of the leaf area is infected. Use higher rate when longer disease control is required.
	Powdery mildew		145 or 290		Apply if more than 5% of stems become infected between full flag emergence to late flowering. When stem rust is the major disease, yield responses are usually optimised by delaying application until full head emergence, and using the higher rate. In severe cases, if a majority of stems are infected prior to full head emergence, apply at 145 mL/ha as soon as possible and if necessary, repeat after 3 weeks when heads are fully emerged.
Wheat Oats	Stem rust				Apply ARYSTA LIFESCENCE TEBCON 430 SC FUNGICIDE onto lime super. Ensure good coverage of all lime super particles.
Onions (Tas only)	White root rot	Before sowing apply ARYSTA LIFESCENCE TEBCON 430 SC FUNGICIDE onto lime super. Ensure good coverage of all lime super particles.	1.45 mL/100 m of row mixed with 145–218 g lime super/100 m of row	–	Apply ARYSTA LIFESCENCE TEBCON 430 SC FUNGICIDE treated lime super when sowing onion seed. Seed and lime super can either be mixed in the same box on the drill or placed in different boxes and sown down the same tube. Apply in a bandwidth of 2cm. Ensure that the correct rate of ARYSTA LIFESCENCE TEBCON 430 SC FUNGICIDE is used otherwise some delay in emergence and reduced stands of seedlings may occur.
Pawpaw	Black spot	Ensure thorough coverage of leaves and fruit.	290	H 3 days	Ensure infected plant material is regularly removed and destroyed to reduce inoculum levels. Spray equipment must be properly calibrated to apply the correct amount of ARYSTA LIFESCENCE TEBCON 430 SC FUNGICIDE. Apply ARYSTA LIFESCENCE TEBCON 430 SC FUNGICIDE at 14 day intervals. Alternate ARYSTA LIFESCENCE TEBCON 430 SC FUNGICIDE with sprays of a protectant fungicide (e.g. mancozeb). Do not apply more than 6 sprays of ARYSTA LIFESCENCE TEBCON 430 SC FUNGICIDE (or any DMI fungicide) on any block in any 12 month period.
Pyrethrum	<i>Sclerotinia sclerotiorum</i>		350	–	Apply twice, in rotation with other control measures, at 7–10 day intervals. Commence at 1 to 2% flowering. Use under direction of pyrethrum advisers.
Ryegrass and fescue seed crops	Leaf rust Stem rust	Apply in at least 100 L of water per ha	290	–	Monitor crops closely and spray at the first signs of disease. Continuing disease pressure or re-infection may require a further application 3–4 weeks later. Ensure thorough coverage and use higher water volumes in dense or advanced crops.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

WITHHOLDING PERIODS:

BANANAS, AVOCADOS: DO NOT HARVEST FOR 1 DAY AFTER APPLICATION.

BEANS, PEAS: DO NOT HARVEST FOR 3 DAYS AFTER APPLICATION AND DO NOT GRAZE OR CUT FOR LIVESTOCK FOR 3 DAYS AFTER APPLICATION.

CEREALS: DO NOT HARVEST FOR 5 WEEKS AFTER APPLICATION AND DO NOT GRAZE OR CUT FOR LIVESTOCK FOR 14 DAYS AFTER APPLICATION.

ONIONS: NOT REQUIRED WHEN USED AS DIRECTED.

PAWPAW: DO NOT HARVEST FOR 3 DAYS AFTER APPLICATION.

PEANUTS: DO NOT HARVEST FOR 21 DAYS AFTER APPLICATION AND DO NOT GRAZE OR CUT FOR LIVESTOCK FOR 21 DAYS AFTER APPLICATION.

H = Harvest, G = Grazing

GENERAL INSTRUCTIONS

FUNGICIDE RESISTANCE WARNING

GROUP 3 FUNGICIDE

ARYSTA LIFESCENCE TEBCON 430 SC FUNGICIDE is a member of the DMI group of fungicides. For fungicide resistance management the product is a Group 3 Fungicide. Some naturally occurring individual fungi resistant to ARYSTA LIFESCENCE TEBCON 430 SC FUNGICIDE and other Group 3 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by ARYSTA LIFESCENCE TEBCON 430 SC FUNGICIDE or other Group 3 fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, Arysta LifeScience Australia Pty Ltd accepts no liability for any losses that may result from the failure of ARYSTA LIFESCENCE TEBCON 430 SC FUNGICIDE to control resistant fungi.

Resistance management recommendations – Peanuts

Apply no more than 3 consecutive sprays of DMI fungicides (e.g. Tebuconazole) before switching to a non-DMI fungicide. Apply no more than 5 DMI sprays per season.

MIXING

Prior to pouring, shake container vigorously, then add the required amount of ARYSTA LIFESCENCE TEBCON to water in the spray vat while stirring or with agitators in motion. Add the required amount of Agridex (peanuts, beans) or water miscible oil (bananas) and mix thoroughly.

APPLICATION

Aircraft should fly as low as possible under the prevailing conditions to minimise drift.

SPECIAL WARNING – BANANAS

There are certain conditions when the surface of recently emerged fruit is particularly prone to marking damage from spray applications. In circumstances where application will be made to very rapidly growing fruit in hot conditions with strong direct light, it is recommended that all emerged bunches be bagged prior to spraying to minimise risk of fruit marking. Never include adjuvants other than water miscible oils with ARYSTA LIFESCENCE TEBCON 430 SC FUNGICIDE sprays. The inclusion of wetting agents is known to cause phototoxicity to young fruit.

FOLIAR DISEASES ON CEREAL CROPS

Do not apply to cereal crops more than once in a season. Treatment will give approximately three weeks disease suppression. Economic response may not be gained by spraying crops past flowering stage. The effects of fungicide application will not be clearly seen for 7–10 days after application. Yield potential: crops with potential yield under 2t/ha are unlikely to give economic responses to a fungicide spray except under conditions of very severe disease. Economic responses are most likely with crops with potential yield of over 3t/ha.

Spray Timings for Stripe Rust Control

Obtain advisory literature from Department of Agriculture for classification of resistant and susceptible varieties.

The following spray program is suggested:

Seedling infections: When approximately 20 out of 100 leaves show signs of infection during tillering or jointing, apply a spray within a week.

Adult infections

Susceptible varieties: When approximately 10 out of 100 leaves show first sign of infection apply a spray within a week.

Moderately Susceptible varieties: When approximately 15–20 out of 100 leaves show first sign of infection apply a spray within a week. Do not delay.

Moderately Resistant and Resistant Varieties: Monitor carefully. If rust appears and spreads, spray within 1 week.

PRECAUTIONS

Re-Entry Period

DO NOT allow entry into treated area until the spray has dried. When prior entry is necessary, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), chemical resistant gloves and footwear. Clothing must be laundered after each day's use.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with chemical or used containers. A spray drift minimisation strategy should be employed at all times when aerially applying sprays. The strategy envisaged is exemplified by the cotton industry's Best Management Practise Manual.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight.

This container can be recycled if it is clean, dry, free of visible residues and has the drumMUSTER logo visible. Triple-rinse container for disposal. Dispose of rinseate or any undiluted chemical according to state legislative requirements. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any drumMUSTER collection or similar container management program site. The cap should not be replaced, but may be taken separately.

For Refillable containers: Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. When opening the container, preparing the spray and using prepared spray wear cotton overalls buttoned to the neck and wrist, washable hat, elbow-length PVC gloves. After use, and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and contaminated clothing.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre. Telephone Australia 13 11 26.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet, which can be obtained from the supplier.

Additional statements required by Globally Harmonised System for classification and labelling of chemicals (GHS) and Safe Work Australia:

Fatal if inhaled. Suspected of damaging fertility or the unborn child.
Do not handle until all safety precautions have been read and understood. Do not breathe dust/fume/gas/mist/vapours/spray. Do not eat, drink or smoke when using this product. Use only outdoors or in a well-ventilated area. Wear respiratory protection.
IF SWALLOWED: Call a POISON CENTER or doctor/physician if you feel unwell. Rinse mouth. IF INHALED: Remove victim to fresh air and keep at rest in a position comfortable for breathing. Immediately call a POISON CENTER or doctor/physician.
IF exposed or concerned: Get medical advice/attention.

MANUFACTURERS WARRANTY AND EXCLUSION OF LIABILITY

Arysta LifeScience Australia Pty Ltd has no control over storage, handling and manner of use of this product. Where this material is not stored, handled or used correctly and in accordance with directions, no express or implied representations or warranties concerning this product (other than non-excludable statutory warranties) will apply. Arysta LifeScience Australia Pty Ltd accepts no responsibility for any loss arising from incorrect storage, handling or use.

In a Transport Emergency
Dial
000
Police or Fire Brigade

<p>UN No. 3082</p>	<p>ENVIRONMENTALLY HAZARDOUS SUBSTANCE, LIQUID, N.O.S. (CONTAINS TEBUCONAZOLE) MARINE POLLUTANT</p>
<p>In a Transport Emergency Dial 000 Police or Fire Brigade</p>	<p>SPECIALIST ADVICE IN EMERGENCY ONLY 1800 033 111 ALL HOURS – AUSTRALIA WIDE</p>
<p>P.G. III</p>	<p>HAZCHEM 3Z</p>

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight.

This container can be recycled if it is clean, dry, free of visible residues and has the drumMUSTER logo visible. Triple-rinse container for disposal. Dispose of rinsate by adding it to the spray tank.

Do not dispose of undiluted chemical on site. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any drumMUSTER collection or similar container management program site. The cap should not be replaced, but may be taken separately.

For Refillable containers: Empty contents fully into application equipment.

Close all valves and return to [point of supply/designated collection point/other specific collection details] for refill or storage.

SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. When opening the container, preparing the spray and using prepared spray wear cotton overalls buttoned to the neck and wrist, washable hat, elbow-length PVC gloves. After use, and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, goggles and contaminated clothing.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre. Telephone Australia 13 11 26.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet, which can be obtained from the supplier.


Additional statements required by Globally Harmonised System for classification and labelling of chemicals (GHS) and Safe Work Australia:

Fatal if inhaled. Suspected of damaging fertility or the unborn child. Do not handle until all safety precautions have been read and understood. Do not breathe dust/ fume/ gas/ mist/ vapours/ spray. Do not eat, drink or smoke when using this product. Use only outdoors or in a well-ventilated area. Wear respiratory protection.

IF SWALLOWED: Call a POISON CENTER or doctor/ physician if you feel unwell. Rinse mouth. IF INHALED: Remove victim to fresh air and keep at rest in a position comfortable for breathing. Immediately call a POISON CENTER or doctor/ physician. IF exposed or concerned: Get medical advice/ attention.

MANUFACTURERS WARRANTY AND EXCLUSION OF LIABILITY

Arysta LifeScience Australia Pty Ltd has no control over storage, handling and manner of use of this product. Where this material is not stored, handled or used correctly and in accordance with directions, no express or implied representations or warranties concerning this product (other than non-excludable statutory warranties) will apply. Arysta LifeScience Australia Pty Ltd accepts no responsibility for any loss arising from incorrect storage, handling or use.



Batch No:

Date of Manufacture:

APVMA Approval No.: 66745/103595